

Athermal Dynamics of Strongly Coupled Stochastic Three-State Oscillators

Bastien Fernandez¹ and Lev S. Tsimring²

¹*Centre de Physique Théorique CNRS, Universités de Marseille I et II et de Toulon-Var, Luminy Case 907, 13288 Marseille CEDEX 09, France*

²*Institute for Nonlinear Science, University of California, San Diego, La Jolla, California 92093-0402, USA*

(Received 8 January 2008; published 25 April 2008)

We study the behavior of globally coupled ensembles of cyclic stochastic three-state units with transition rates from $i - 1$ to i proportional to the number of units in state i . Contrary to mean-field theory predictions, numerical simulations show significant stochastic oscillations for sufficiently large coupling strength. The order parameter characterizing units synchrony increases monotonically with coupling while the coherence of oscillations has a maximum at a certain coupling strength. We find the exact formulas for the stationary probability distribution and the order parameter.

DOI: [10.1103/PhysRevLett.100.165705](https://doi.org/10.1103/PhysRevLett.100.165705)

PACS numbers: 64.60.Ht, 05.45.Xt, 89.75.-k

Interacting stochastic systems emerge in a variety of physical and biological contexts, from arrays of Josephson junctions [1] and lasers [2] to neural networks [3,4] and gene regulatory networks [5,6]. While the dynamics of individual elements can be rather complicated and nongeneric, large ensembles of coupled units often exhibit universal behavior. Therefore, studies of canonical models with simple individual dynamics and interaction rules have proven very useful for understanding the behavior of specific systems. Well-known examples of such canonical systems are the Desai-Zwanzig model [7] of coupled bistable systems and the Kuramoto model [8] of coupled phase oscillators. The Kuramoto model and its many variations and generalizations (see [9]) have been very successful in describing the transition to coherent oscillations in ensembles of coupled phase oscillators. Stochastic dynamics of individual elements in such models are described by coupled nonlinear Langevin equations. In the thermodynamic limit, they can be reduced to low-dimensional deterministic equations for the mean field or the order parameter characterizing global behavior of large systems.

A simpler way of describing interacting stochastic systems incorporates stochastic elements with a discrete set of states with certain transition rates. It is most often done for bistable systems which are replaced by two-state stochastic systems with suitably chosen transition rates. For example, array-enhanced stochastic resonance has been studied in a system of globally coupled two-state systems [10]. A transition to regular oscillations in an ensemble of two-state systems coupled through a delayed mean field was studied in [11].

Recently, Prager *et al.* [12] introduced a globally coupled three-state stochastic “oscillators” with unidirectional ($1 \rightarrow 2 \rightarrow 3 \rightarrow 1$) transitions as a paradigmatic model of noise-driven excitable systems. This model is simple enough to make analytical and large-scale numerical studies of large systems feasible [12,13]. An important property of globally coupled systems is their behavior in the thermodynamic limit, when the number of units ap-

proaches infinity. According to the standard mean-field description, this behavior depends on the specific form of coupling among oscillators: in certain cases [13] the ensemble may exhibit a transition to sustained oscillations (Hopf bifurcation) for strong enough coupling, and in some others [12] this transition is absent.

In this Letter we focus on the seemingly “less interesting” situation when a globally coupled ensemble of three-state systems *does not* exhibit a Hopf bifurcation in the thermodynamic limit. However, we find that in any finite-size system, quasiregular oscillations of the mean field are present. We introduce the coherence parameter which characterizes regularity of mean-field oscillations, and the order parameter which characterizes the degree of synchrony among the oscillators. We show that while the order parameter increases monotonically with the coupling strength, the coherence parameter has a maximum at a certain intermediate coupling strength. The simplicity and a high degree of symmetry in the system under study allow us to find the statistical properties of the finite ensemble analytically; however, we expect similar behavior in ensembles of more general cyclic stochastic systems at strong enough coupling which do not exhibit Hopf bifurcation in the thermodynamic limit.

A single stochastic three-state unit with unidirectional transitions is schematically shown in the inset to Fig. 1(b). We assume that in an isolated unit all three transitions from state i ($i = 1, 2, 3$) to state $i + 1 \pmod{3}$ are Markovian with identical rate a [14]. Without loss of generality we take $a = 1$. Statistical properties of this system have been investigated in Ref. [12]. The cyclic behavior of a single oscillator is characterized by the mean time T of an oscillator to return to the initial state after an excursion through the other two states. Since the mean time of switching from state i to state $i + 1$ is $T_s = 1$, we get $T = 3T_s = 3$. The probability for a system to be in state $i = 1, 2, 3$ at time t is given by the continuous-time master equation

$$\dot{P}_i = -P_i + P_{i-1}, \quad i = 1, 2, 3 \pmod{3}. \quad (1)$$




FIG. 1 (color online). Time series of the instantaneous occupation numbers oscillators n_1, n_2, n_3 in states 1, 2, 3, respectively, for $N = 1000$, and different values of the coupling parameter: (a) $b = 0$, (b) $b = 0.1$ (Inset: Transition diagram in a single unit); (c) $b = 1$, (d) $b = 10$.

This master equation has a fixed point $P_1^s = P_2^s = P_3^s = 1/3$ corresponding to equipartition among the three states, and three eigenvalues $\lambda_k = -1 + e^{2i\pi k/3}$, $k = 0, 1, 2$. The first eigenvalue ($k = 0$) corresponds to the conservation of the total probability, and the other two describe equilibration of the probability distribution among the three states. Imaginary part of these eigenvalues implies that there are decaying periodic oscillations of deviations from equipartition with the mean frequency $\omega = \sqrt{3}/2$.

Globally coupled three-state oscillators.—Now let us consider an ensemble of N identical three-state oscillators. The specific mechanism of coupling is the following. We assume that the probability $\pi_{i,i+1}$ of switching of an oscillator from a state i to state $i + 1$ at time t is linearly proportional to the number of oscillators $n_{i+1}(t)$ already at state $i + 1$ at time t , with the proportionality constant b (which we call *coupling coefficient*), $\pi_{i,i+1}(t) = 1 + bn_{i+1}(t)$. This type of coupling is reminiscent of the autocatalytic transitions in gene regulatory circuits when multiple copies of a single gene are present in the cell.

Since this model is Markovian, it can be efficiently simulated using Gillespie algorithm [15]. Figure 1 shows sample stochastic trajectories for the occupation number of oscillators in states 1, 2, and 3 as a function of time for different values of coupling parameter b for $N = 1000$. The initial condition for all cases is $n_1(0) = N$, $n_{2,3}(0) = 0$. For $b = 0$, the population slowly drifts toward an equilibrium state with $n_1 = n_2 = n_3 = N/3$ with $O(N^{1/2})$ stochastic fluctuations. For even very small nonzero $b \ll 1$, noisy oscillations around the mean become visible. As b grows, the period becomes shorter, and the amplitude of oscillations grows until for $b \sim 1$ it approaches N ; i.e., almost all oscillators are simultaneously in the same state. At large $b \gg 1$, the system exhibits switching behavior resembling the dynamics of a single oscillator. It is easy to understand: at very large b , once a single oscillator makes a transition from state i to state $i + 1$ (which occurs with rate N), all other oscillators quickly follow. So indeed in the limit $b \gg 1$ the dynamics of the ensemble becomes equivalent to the dynamics of a single oscillator with a

rescaled transition rate N ; a result confirmed by analytical calculations [16].

We computed the power spectrum of the time series of the occupation numbers and determined the central frequency ω and the half-width $\Delta\omega$ of the spectral peak. We call the ratio $CP = \omega/\Delta\omega$ the coherence parameter. Figure 2(a) shows ω , $\Delta\omega$, and CP as functions of the coupling parameter b . Both ω and $\Delta\omega$ increase with b ; however, the coherence parameter has a distinct maximum at a certain b which decreases with N ; see Fig. 2(a). Thus, we observe a manifestation of the coherence resonance [17]; however, the difference is that the maximum appears not at a certain noise strength but a certain value of the coupling.

This coherence resonance should not be confused with *synchronization* among the oscillators. The degree of synchronization can be characterized by the order parameter

$$R = \left\langle N^{-1} \left| \sum_{j=1}^N e^{i\phi_j} \right|^2 \right\rangle, \quad (2)$$

with discrete phases of oscillators $\phi_j = 2\pi k/3$, $k = 1, 2, 3$. This order parameter was introduced by [13] by analogy with coupled continuum phase oscillators in the Kuramoto model [8]. The order parameter is zero when all oscillators are equally distributed among the three states ($n_1 = n_2 = n_3 = N/3$), and it approaches unity if all oscillators are perfectly synchronized, so all of them are simultaneously in the same state. As expected, the order parameter increases monotonically with b ; see Fig. 2(b). As Fig. 2(b) shows, the order parameter for a given b is *independent* of the number of oscillators in population N .

Mean-field approximation.—In the thermodynamic limit $N \rightarrow \infty$ of globally coupled oscillators, the mean-field equations for the deterministic “concentrations” of oscillators $x_i = n_i/N$ are given by

$$\begin{aligned} \dot{x}_i &= x_{i-1}(1 + bNx_i) - x_i(1 + bNx_{i+1}), \\ i &= 1, 2, 3(\text{mod}3). \end{aligned} \quad (3)$$

From the symmetry of the underlying dynamics it immediately follows that the fixed point of this system is $x_1 = x_2 = x_3 = 1/3$. The associated two complex eigenvalues (the first eigenvalue is 0 as before because of the conser-




FIG. 2 (color online). Gillespie simulations of coupled stochastic oscillators: coherence parameter $CP = \Delta\omega/\omega$ (a) and order parameter R (b) vs. b for $N = 10^2, 10^3, 10^4$. Solid line in (b) is (8). Inset: frequency ω and bandwidth $\Delta\omega$ vs b .

vation law) $\lambda_{2,3} = -\frac{3}{2} \pm i\sqrt{3}(\frac{1}{2} + \frac{bN}{3})$ always have negative real part, which indicates the absence of Hopf bifurcation at any coupling b . This is in contrast with the model by Wood *et al.* [13], in which a different form of coupling was proposed (the transition rate from i to $i+1$ was an exponential function of the linear combination of $n_{i\pm 1}, n_i$), for which the mean field had a Hopf bifurcation at large enough coupling strength [18].

Stochastic description.—The full description of the stochastic properties of the system is given by the master equation for the probability $p(n_1, n_2, n_3)$ that at time t there are n_1 oscillators at state 1, n_2 at state 2, and n_3 at state 3 (obviously, $n_1 + n_2 + n_3 = N$),

$$\begin{aligned} \dot{p}(n_1, n_2, n_3) = & (n_1 + 1)[1 + b(n_2 - 1)] \\ & \times p(n_1 + 1, n_2 - 1, n_3) + (n_2 + 1) \\ & \times [1 + b(n_3 - 1)]p(n_1, n_2 + 1, n_3 - 1) \\ & + (n_3 + 1)[1 + b(n_1 - 1)] \\ & \times p(n_1 - 1, n_2, n_3 + 1) \\ & - [N + b(n_1 n_2 + n_2 n_3 + n_3 n_1)] \\ & \times p(n_1, n_2, n_3). \end{aligned} \quad (4)$$

The total number of states (n_1, n_2, n_3) in this system is $(N+2)(N+1)/2$. It is convenient to depict the state space as a triangular grid; see Fig. 3(a).

According to the Frobenius-Perron theorem, the master equation has a unique stationary solution. We were able to find the exact solution in a closed form,

$$p_s(n_1, n_2, n_3) = C_b \frac{G(b, n_1)G(b, n_2)G(b, n_3)}{n_1!n_2!n_3!}, \quad (5)$$

where $G(x, n) = \prod_{k=0}^{n-1} (1 + kx)$ and C_b is the normalization constant, which can be verified by direct substitution. Expression (5) shows that the stationary solution is not only invariant with respect to cyclic permutations but also with respect to any permutation of coordinates (n_1, n_2, n_3) . This property is remarkable because the equation itself does not possess this symmetry.

It is easy to check by direct substitution that (5) satisfies the relation

$$\begin{aligned} (n_3 + 1)[1 + b(n_1 - 1)]p_s(n_1 - 1, n_2, n_3 + 1) \\ = n_1(1 + bn_3)p_s(n_1, n_2, n_3) \end{aligned} \quad (6)$$

and two other relations obtained from (6) by cyclic permutations (adding these 3 relations and using the rotation symmetry gives the stationary master equation Eq. (4) without the left-hand side).

It is straightforward to obtain convexity properties of the stationary solution (5). Indeed, according to (6), we have $p_s(n_1 - 1, n_2, n_3 + 1) < p_s(n_1, n_2, n_3)$ iff $0 < (n_3 + 1 - n_1)(1 - b)$. If $b < 1$ ($b > 1$) the probability increases (decreases) when one moves one step to the right in the left part of the triangular lattice and vice versa [19]. Combining it with the rotation symmetry, we conclude that the distri-





FIG. 3 (color online). (a) State space of the three-oscillator system $N = 3$; Vertices correspond to distinct states (n_1, n_2, n_3) of the system, and arrows indicate transitions among the states. Expressions at the arrows show the corresponding transition rates. (b)–(e) Stationary probability distributions for $N = 14$ and $b = 0.2$ (b), $b = 0.8$ (c), $b = 1$ (d), $b = 2$ (e).

bution is convex with a maximum in the center when $b < 1$ and concave with a maximum in the corners for $b > 1$; for $b = 1$ the distribution is flat.

For zero coupling ($b = 0$), the stationary distribution is trinomial

$$p_s(n_1, n_2, n_3) = \frac{N!}{3^N n_1! n_2! n_3!},$$

which, of course, could be deduced directly since oscillators are independent and in the long term limit they become uniformly distributed among the three states. The most probable state is in the middle of the triangle ($n_1 = n_2 = n_3 = N/3$) and the least probable states are in the corners $(N, 0, 0)$, $(0, N, 0)$, $(0, 0, N)$.

For large b , the stationary distribution is highly localized at the corners [Fig. 3(e)]. The dynamics close to equilibrium can be approximated by the probability flow around the edges of the triangle, ignoring the influence of the inner nodes. This simplification allows us to compute the non-

zero eigenvalues of the full system in the first order in $1/b$ [16]. For large b these eigenvalues are $\lambda_k N + O(1/b^2)$, $k = 1, 2$, where λ_k are the eigenvalues of the single oscillator. Note that the equilibration rate $\text{Re}(-\lambda_{1,2}N)$ is independent of b .

For large n_1 , n_2 and n_3 one can use Stirling formula to find an asymptotic expression for the distribution (5),

$$p_s(n_1, n_2, n_3) = C_b(n_1 n_2 n_3)^{1/b-1} \quad (7)$$

with $C_b = \Gamma(3/b)\Gamma^{-3}(1/b)N^{1-3/b}$. We can use this expression, replace summation by integration and compute the order parameter for large N explicitly [20]. This straightforward calculation results in a surprisingly simple formula

$$R = \frac{b}{b+3}. \quad (8)$$

This formula agrees very well with direct Gillespie simulations [Fig. 2(b)]. Note that the order parameter is independent of N (at least for large N). For arbitrary N , the order parameter can be computed from the stationary distributions at zero coupling and large coupling, respectively. For zero coupling, we get $R = 1/N$ and for large coupling, we have $R \rightarrow 1$.

Discussion.—Our analysis indicates that the large systems of globally coupled three-state units exhibit significant stochastic oscillations, and while the frequency of these oscillations scales as N , their temporal coherence reaches maximum at a finite $b \sim 1$ independent of the number of oscillators. This result is counterintuitive, since the mean-field theory predicts no sustained oscillations in the thermodynamic limit $N \rightarrow \infty$. The origin of this apparent contradiction is that for large coupling strength, nearly all oscillators are clustered in a single state, and other states are nearly empty, thus giving rise to strong stochasticity. For large b , all oscillators are strongly correlated: as soon as the first one makes a transition, others very quickly follow. This leads to the noisy but correlated and thus athermal behavior of the globally coupled system at large b . Of course, in any real system the transition rate should saturate as $N \rightarrow \infty$. Then eventually the thermal behavior of the system would be recovered; however, in the intermediate scaling regime of finite N the dynamics described here can be observed.

Our analytical results are easily generalizable to the case of arbitrary d -state oscillators with any $d > 1$. The order parameter for d -oscillators is simply $R_d = b/(b+d)$. For all $d > 2$, the coherence parameter reaches maximum at $b \sim 1$. However, two-state systems ($d = 2$) are qualitatively different since for those the detailed balance condition is satisfied, and the probability current is zero. In this case, the quasiregular oscillations are completely absent, and the coherence parameter remains small $O(1)$ for any b (see [16] for more details).

The authors are indebted to K. Lindenberg for an illuminating discussion. L. T. is grateful to CNRS and to University of Aix-Marseille I for hospitality and support. This work was partially supported by ACI IMPBio (BF) and MURI ONR No. N00014-07-1-0741 and NIH Grant No. GM69811-01 (L. T.).

-
- [1] P. Hadley, M. R. Beasley, and K. Wiesenfeld, *Phys. Rev. B* **38**, 8712 (1988).
 - [2] H. Bruesselbach, D. C. Jones, M. S. Mangir, M. Minden, and J. L. Rogers, *Opt. Lett.* **30**, 1339 (2005).
 - [3] M. Abeles, *Corticonics: Neural Circuits of the Cerebral Cortex* (Cambridge University Press, Cambridge, England, 1991).
 - [4] M. Tsodyks, K. Pawelzik, and H. Makram, *Neural Comput.* **10**, 821 (1998).
 - [5] J. Hasty, J. Pradines, M. Dolnik, and J. Collins, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 2075 (2000).
 - [6] J. Paulsson, *Nature (London)* **427**, 415 (2004).
 - [7] R. Desai and R. Zwanzig, *J. Stat. Phys.* **19**, 1 (1978).
 - [8] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence* (Springer-Verlag, Berlin, 1984).
 - [9] J. Acebrón, L. Bonilla, C. Pérez Vicente, F. Ritort, and R. Spigler, *Rev. Mod. Phys.* **77**, 137 (2005).
 - [10] J. F. Lindner, M. Bennett, and K. Wiesenfeld, *Phys. Rev. E* **73**, 031107 (2006).
 - [11] D. Huber and L. S. Tsimring, *Phys. Rev. Lett.* **91**, 260601 (2003).
 - [12] T. Prager, B. Naundorf, and L. Schimansky-Geier, *Physica (Amsterdam)* **325A**, 176 (2003).
 - [13] K. Wood, C. Van den Broeck, R. Kawai, and K. Lindenberg, *Phys. Rev. Lett.* **96**, 145701 (2006); *Phys. Rev. E* **74**, 031113 (2006); **75**, 041107 (2007); *Phys. Rev. E* **76**, 041132 (2007).
 - [14] This assumption is made merely for analytical tractability, the system behavior remains qualitatively the same for arbitrary nonequal transition rates.
 - [15] D. Gillespie, *J. Phys. Chem.* **81**, 2340 (1977).
 - [16] See EPAPS Document No. E-PRLTAO-100-022817 for details of the rigorous analytic computation of the main eigenvalues of the full system in the large coupling limit and the generalization of the theory for the order parameter to d -state oscillators with arbitrary $d > 1$. For more information on EPAPS, see <http://www.aip.org/pubservs/epaps.html>.
 - [17] A. S. Pikovsky and J. Kurths, *Phys. Rev. Lett.* **78**, 775 (1997).
 - [18] For a general nonlinear transition probability $\pi_{i,i+1} = \Pi(x_{i+1})$, the real part of the eigenvalues is given by $[\Pi'(1/3) - 3\Pi(1/3)]/2$, so the Hopf bifurcation occurs only if the relative slope of the coupling function $\Pi'(1/3)/\Pi(1/3)$ can exceed 3.
 - [19] The left part of the lattice corresponds to points for which $n_1 \leq n_3 - 1$.
 - [20] When N is large, the states (n_1, n_2, n_3) close to the edges, where the expression (7) is not valid, give a little contribution to the order parameter.